

1. Mild EPH - Gestosis (Score less than 4):

systolic BP. up to	159 mmHg	(1)
and/or diastolic BP up to	99 mmHg	(1)
and/or oedema up to leg		(1)
and/or proteinuria up to	0.5 gm/L	(1)

2. Severe EPH - Gestosis (Score between 4 & 8)

systolic BP	160-179	(2)
and/or diastolic BP	100-109	(2)
and/or oedema	legs / abdomen	(2)
and/or proteinuria	< 0,5 gn/L	(2)

3. Imminent Eclampsia (IE)
Severe EPH Gestosis
plus mental, visual, auditory, cloudiness & hallucinations
AND/OR epigastric pain & vomiting
AND/OR hyperreflexia, oliguria & anuria

4. Fulminating Eclampsia (FE)
Symptoms & signs proceeding from normal to severe EPH - Gestosis
or imminent Eclampsia within a period of 12 hours or less.

5. Eclampsia (CE)
EPH - Gestosis with convulsive fits

Management guidelines.

1. Span of treatment to see if the condition is responding or not.
 - a. Mild : 12 to 15 days
 - b. Severe: 3 to 5 days
 - c. Imminent and fulminating : 6 to 12 hours

2. Location of treatment. Place recommended for management of the patient and for her observation.
 - a. Mild : at home
 - b. Severe: at antenatal ward
 - c. Imminent & Fulminating : In ICU (special IUC equipped for management of EPH - Gestosis)

3. Two solid lines of treatment (sure to be effective)