

Gestosis

Laboratory methods of examination for immunoincompatibility married couples. Out and during pregnancy.

Goal: for detection of the causes and correction thereof.

- < Immunoincompatibility of couple (ABO, RH, Kell, Duffy, Kidd, MNSs, HLA, A,B, C, DR (Loci). Index hystocompatibility (HLA) between couples.
- < Detection antierythrocyte (complete and incomplete) and antileucocyte (anti-HLA) antibodies.
- < TORCH-STDs infections
- < Immunogram- index immunoregulatority Th/Ts, Immunoglobulin levels:Ig G, Ig A, Ig M.

Ph.D. Immuno-serologist Tsisana Chagiashvili.

Institute of Hematology and blood transfusion. Tbilisi. Georgia.

Ph.D. Tsisana Chagiashvili, Immuno-serologist, Georgian Institute of Haematology and blood transfusion. Scientific worker. (Tbilisi, Georgia).

Email: tschagia@yahoo.com.

IMMUNOLOGICAL COMPATIBILITY OF MARRIED COUPLES IN ETHIOPATHOGENESIS OF GESTOSIS OR OUT OF PREGNANCY

LABORATORY METHODS:

- < Serological tests:
Blood typing ABO, RH, MNSs, Kell, DYffy, Kidd, Blood systems antigens (tube test).
- < HLA typing A, B, C, DR Loci antigens (Terasaki test). Index hystocompatibility
- < TORCH-STD infections (ELISA test)
- < Immunological test: Immuno-fluerecence CD3, CD4, CD8, CD20. Index Immunoregulatority Th/Ts
- < Immunoglobulin levels (Ig G, Ig M, Ig A)- Manchini test.